

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

MR 73

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/54, 15/10, 9/12, C12Q 1/48, A61K 31/70		A1	(11) International Publication Number: WO 95/13383
			(43) International Publication Date: 18 May 1995 (18.05.95)
(21) International Application Number: PCT/US94/13130		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 10 November 1994 (10.11.94)		Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(30) Priority Data: 08/153,051 12 November 1993 (12.11.93) US 08/151,477 12 November 1993 (12.11.93) US			
(71) Applicants: GERON CORPORATION [US/US]; 200 Constitution Drive, Menlo Park, CA 94025 (US). BOARD OF REGENTS, THE UNIVERSITY OF TEXAS SYSTEM [US/US]; 201 West Seventh Street, Austin, TX 78701 (US).			
(72) Inventors: SHAY, Jerry; 5060 Horseshoe Trail, Dallas, TX 75209 (US). WEST, Michael, David; 15 Wakefield Court, Belmont, CA 94002 (US). WRIGHT, Woodring, E.; 2300 Country Green Lane, Arlington, TX 76011 (US).			
(74) Agents: WARBURG, Richard, J. et al.; Lyon & Lyon, First Interstate World Center, Suite 4700, 633 West Fifth Street, Los Angeles, CA 90017-2066 (US).			
(54) Title: METHODS AND REAGENTS FOR LENGTHENING TELOMERES			
(57) Abstract Method and compositions for increasing telomere length in normal cells can be used to increase the proliferative capacity of cells and to delay the onset of cellular senescence.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

TITLE

Methods and Reagents for Lengthening Telomeres

5

10

15

Background of the InventionField of the Invention

This invention relates to methods and reagents for extending the telomeres and therefore the proliferative capacity of cells. The invention therefore relates to the fields of chemistry, pharmacology, biology, molecular biology, and medical therapeutic and diagnostic technology.

20

Description of Related Art

Normal human somatic cells (i.e., fibroblasts, endothelial, and epithelial cells) display a finite replicative capacity of 50-100 population doublings characterized by a cessation of proliferation in spite of the presence of growth factors. This cessation of replication in vitro is variously referred to as cellular senescence or cellular aging. See, Goldstein, 1990, Science 249:1129, and Hayflick and Moorehead, 1961, Exp. Cell Res. 25:585. The replicative life span of cells is inversely proportional to the in vivo age of the donor, so cellular senescence is believed to play an important role in aging in vivo. See Martin et al., 1979, Lab. Invest. 23:86, and Schneider and Mitsui, 1976, Proc. Natl. Acad. Sci. USA 73:3584.

30

35

Cellular immortalization (the acquisition of unlimited replicative capacity) may be thought of as an abnormal escape

from cellular senescence. See Shay *et al.*, 1991, Exp. Cell Res. 196:33. Normal human somatic cells appear to be mortal and to have finite replicative potential. In contrast, germline and malignant tumor cells appear to be immortal and to have indefinite proliferative potential. Human cells cultured *in vitro* appear to require the aid of transforming viral oncoproteins to become immortal, and even then, the frequency of immortalization is 10^{-6} to 10^{-7} . See Shay and Wright, 1989, Exp. Cell Res. 184:109. Cells obtained from murine sources immortalize at a relatively high frequency without the aid of transforming oncoproteins. A variety of hypotheses have been advanced over the years to explain the causes of cellular senescence.

Shay *et al.*, 1992, Experimental Gerontology 27:477, and Shay *et al.*, 1991, Exp. Cell Res. 196:33, describe a two-stage model for human cell mortality to explain the ability of Simian Virus 40 (SV40) T-antigen to immortalize human cells. The mortality stage 1 mechanism (M1) is a mechanism by which cells cease to proliferate after a certain number of population doublings, and the biological molecules that carry out this mechanism appear to be the target of certain tumor virus proteins. An independent mortality stage 2 mechanism (M2) produces crisis in cells that have bypassed M1, with crisis being typified by severe chromosomal abnormalities and ultimately cell death. The M2 mechanism thus prevents tumor viruses from directly immortalizing human cells. The papers noted above describe experiments in which T-antigen expression was driven by a mouse mammary tumor virus promoter to cause reversible immortalization of cells. SV40 T-antigen extends the replicative life span of human fibroblasts by an additional 40-60%. The M1 mechanism is overcome by T-antigen, perhaps by binding to various cellular proteins or by inducing new activities to repress the M1 mortality mechanism. The M2 mechanism then causes cessation of proliferation, even though the M1 mechanism is blocked. Immortality is achieved only when the M2 mortality mechanism is also disrupted. The M2 mechanism appears to cause a dominant repression of the immortal phenotype,

because hybrids between mortal and immortal human cells are generally mortal. See Pereira-Smith and Smith, 1983, Science 221: 964-966. As described above, such hybrids between murine cells immortalize at a much higher frequency, so it is relatively easy to obtain immortal hybrids between murine lymphocyte and myeloma cells to obtain immortal antibody-producing hybridomas, for instance. However, such hybrids are not immortal when produced with human cells.

The finite replicative capacity of cells may reflect the work of a "clock" linked to DNA synthesis at the telomeres (ends) of chromosomes. Thus, Harley *et al.*, 1990, Nature 345:458, state that the amount and length of telomeric DNA in human fibroblasts decreases as a function of serial passage during aging *in vitro*. Harley, 1991, Mutation Research 256:271, describes that telomeres of human somatic cells act as a mitotic clock, shortening with age both *in vitro* and *in vivo* in a replication dependent manner and proposes a hypothesis for human cell aging and transformation in which telomeres and telomerase, a ribonucleoprotein polymerase involved in telomeric DNA synthesis, play a causal role in cell senescence and cancer.

PCT patent publication No. 93/23572 describes therapeutic and diagnostic methods relating to telomerase and telomere length. The publication describes oligonucleotide reagents that either reduce loss of telomere length during passage of cells *in vitro* or increase telomere length of immortal cells *in vitro*.

There remains a need for reagents that can increase telomere length in mortal cells and methods for extending the proliferative capacity of mortal cells and hybrids between mortal and immortal human cells. The present invention meets these and other needs, as described below, first in summary fashion and then in detail with examples.

Summary of the Invention

The present invention provides a method for increasing the proliferative capacity of normal cells, which method

comprises culturing or cultivating said cells in the presence of a therapeutically effective amount of an oligonucleotide substrate for telomerase under conditions such that said oligonucleotide substrate enters said cells and acts to lengthen telomeric DNA of said cells. The therapeutic method of this invention postpones cellular senescence in cells, and is especially useful for cells expressing telomerase and for hybrids between immortal and mortal cells. The method can also be used to reduce the level of chromosomal fusions and other chromosomal aberrations by increasing the telomere repeat length in cells.

While the cells treated by the present method can be cultured or cultivated in vivo during administration of the oligonucleotide, the invention has particular application to the cultivation of cells ex vivo, and provides especially important benefits to therapeutic methods in which cells are cultured ex vivo and then reintroduced to a host. Especially preferred examples of such cells include the CD4⁺ cells and the CD8⁺ cells of HIV-infected individuals and immune cells of aged people. The method is also preferred for increasing the replicative capacity of hybrids between immortal and mortal human cells, such as hybrids between human B-lymphocytes and myeloma cells, to increase the replicative capacity of antibody producing human hybridomas.

While the method of the invention is applicable to any normal cell type, the method can be practiced using normal cells that express a low level of telomerase activity. For purposes of the present invention, the term "normal" refers to cells other than tumor cells, cancer cells, or transformed cells. Especially preferred cells for use in the present method include embryonic, fetal, neonatal, and adult stem cells of any organ, and adult pluripotent hematopoietic stem cells. The method can also be applied to immortalized or transformed cells, such as myeloma cells, to increase telomere length prior to hybridization with a mortal cell.

The reagents of the invention include any oligonucleotide substrate for telomerase. The oligonucleotides of the invention can be composed of naturally occurring nucleotides

or synthetic nucleotides or synthetic nucleotide analogs or combinations of any of the same. Typically, such oligonucleotides will be at least 12 nucleotides in length or longer.

5 These and other aspects of the invention are described in more detail below, beginning with a brief description of the accompanying drawings.

Brief Description of the Drawings

10 Figure 1 shows the results of telomere length analysis (DNA was isolated from the cells, digested with a restriction enzyme that does not cleave telomeric repeat sequences, size-separated by gel electrophoresis, and probed with radioactively labeled probes comprising telomeric repeat
15 sequences) of IDH4 cells treated according to the method of the invention as well as control cells. The telomeres of IDH4 cells elongate by several kbp after 90 days of thrice weekly treatment in culture with 30 μ M GTR (for "G-rich Telomeric Repeat") oligonucleotide (5'-TTAGGGTTAGGG-3'). Lanes 1, 3,
20 5, and 7 show the results for control, untreated cells. Lane 2 shows the results for cells treated for 90 days with 30 μ M GTR oligonucleotide. Lane 4 shows the results for cells treated for 90 days with 30 μ M GTR oligonucleotide and then cultured for 30 days in the absence of added GTR
25 oligonucleotide. Lane 6 shows the results for cells treated for 90 days with 30 μ M GTR oligonucleotide and then cultured for 60 days in the absence of added GTR oligonucleotide. Lane 8 shows the results for cells treated for 90 days with 30 μ M GTR oligonucleotide and then cultured for 90 days in the
30 absence of added GTR oligonucleotide.

35 Figure 2 shows the results of telomere length analysis of IDH4 cells treated according to the method of the invention as well as control cells and shows that a progressive (up to 30 μ M) dose response is observed with increasing concentrations of the GTR oligonucleotide. Lane 1 shows the results for untreated control cells. Lane 2 shows the results for cells treated for 90 days with 10 μ M GTR oligonucleotide. Lane 3 shows the results for cells treated for 90 days with 30 μ M

6

GTR oligonucleotide. Lane 4 shows the results for cells treated for 90 days with 100 μ M GTR oligonucleotide.

Figure 3 shows the results of telomere length analysis of a variety of different cells treated according to the method of the invention as well as control cells and shows that the effect of the GTR oligonucleotide is not restricted to a few immortal cell lines, because all of the cell lines tested responded to GTR oligonucleotide treatment by elongating their telomeres. In addition to SW620 colon adenocarcinoma cells (lane 1, control, and lane 2, 30 μ M GTR-treated cells after 20 population doublings), CAK1 renal carcinoma cells increase the length of their telomeres approximately the same amount as the SW620 GTR-treated cells (lane 3, control, and lane 4, GTR-treated cells) under the same conditions. In addition, DU145 prostatic adenocarcinoma cells and COLO205 colon carcinoma cells, which have slightly longer telomeres in control (untreated) cells, also elongate their telomeres in the presence of GTR oligonucleotide (lanes 5 and 7, controls, and lanes 6 and 8, GTR-treated, respectively) under the same conditions.

Description of the Preferred Embodiments

All normal diploid vertebrate cells have a limited capacity to proliferate, a phenomenon that has come to be known as the Hayflick limit or replicative senescence. In human fibroblasts, this limit occurs after 50-100 population doublings, after which the cells remain in a viable but non-dividing senescent state for many months. This contrasts to the behavior of most cancer cells, which have escaped from the controls limiting their proliferative capacity and are effectively immortal.

One hypothesis to explain the cause of cellular senescence concerns the role of the distal ends of chromosomes called telomeres. DNA polymerase synthesizes DNA in a 5'-to-3' direction and requires a primer to initiate synthesis. Because of the mechanism of action of DNA polymerase, the "lagging strand" is not replicated to the very end of a linear chromosome, as there is no means to initiate synthesis at the

end of this strand. The chromosome is thus shortened with every cell division.

5 Telomeres comprise long stretches of duplex nucleic acid composed of a repeating sequence, called the "telomeric repeat," which in humans is 5'-TTAGGG-3'. In addition, the G-rich strand of the telomere is longer than the C-rich strand; consequently, telomeres also comprise a single-stranded region composed entirely of telomeric G-rich repeat sequences. The enzyme telomerase is believed to mediate the
10 synthesis of at least the terminal portion of the G-rich strand of the telomere. The cellular senescence hypothesis is that somatic cells lack telomerase and so lack the ability to replicate the very ends of DNA molecules. This results in a progressive shortening of the ends of the chromosomes with
15 each cell division until a point at which this progressive loss signals functional changes, at which time the cell loses the capacity to proliferate.

20 When a cell loses in this manner the capacity to proliferate, the cell is senescent. Thus, the term "senescence" refers to the loss of ability of a cell to replicate in the presence of normally appropriate mitogenic signals. Senescence is also typically accompanied by a change in expression patterns of one or more genes. For instance, senescence in some cells is accompanied by an increase in the
25 expression of degradative enzymes, such as collagenase. The term senescence does not include quiescent cells that can be induced to re-enter the cell cycle under appropriate conditions.

30 Thus, the length of a telomere is directly correlated to the life span of an individual cell. Certain populations of cells may lose telomere length at a greater rate than other cells within an individual, and those cells with shortened telomeres may thus become age-limiting within an individual organism. In such cases, important benefits can be achieved by slowing
35 the loss of telomere sequences, which can be achieved by increasing the length of telomeric DNA.

 The present invention provides a method for increasing the length of telomeres in normal cells and in hybrids

8

between normal and immortal cells, thus extending the ability of a cell or hybrid to replicate and increasing the proliferative capacity of the treated cell. In this method, a telomere-extending amount of an agent that increases telomere length within the cell is provided to the cell. Specifically, the telomeres within a particular cell population can be lengthened by provision of an oligonucleotide that acts to lengthen telomeres in cell culture or cultivation, thus increasing the number of cell divisions that can occur before a cell becomes senescent. By virtue of the present method, one can lengthen telomeres by providing a nucleic acid, e.g., DNA or RNA (including modified forms), that can act as a telomerase substrate to the cells. Such nucleic acids typically include at least 2 to 3 telomeric repeat sequences, which in humans is 5'-TTAGGG-3'.

This method is useful for the treatment of individuals not suffering from any particular condition but in which one or more cell types are limiting in that individual. The method thus can be practiced to delay the onset of cell senescence, which, as noted above, is characterized by the inability of that cell to replicate further in an individual. When practiced in this way, the method is a disease prevention method. The present invention also provides important new methods for treating disease, especially diseases characterized by an increased rate of proliferation of one or more particular groups of cells. An increase in the rate of proliferation increases the rate at which telomeric DNA is lost and the rate at which cells become senescent. The present method can be used to prevent the detrimental effects of early senescence of the cells due to increased cellular turnover associated with the disease.

The present invention also has application in tissue culture techniques to delay the onset of cellular senescence, whether for therapeutic or other purposes, such as manufacturing and research purposes. For instance, many cell-based therapies require the clonal expansion of cells for later reintroduction into a patient. During the expansion process, however, the cells continue to approach the Hayflick

limit, so the cells reintroduced to the patient may be senescent or close (i.e., within a few divisions) to a senescent state. This invention allows the expansion of cells in the presence of telomere lengthening agents and so provides a means to avoid the problems associated with cell senescence. In addition, the present invention allows normal cells to be cultivated for extended doublings in vitro.

Although the present invention has a wide variety of applications, the invention has application in an area of critical importance today: cell therapy. Cell therapy involves the isolation of healthy human cells, the expansion of those cells ex vivo, and the reinfusion of the expanded cells into a patient. Cell therapy has application in the treatment of cancer and organ transplantation and many other disease states or conditions. For instance, bone marrow therapy takes advantage of the fact that bone marrow, the major organ of the immune system, is responsible for production of various cells in the blood from hematopoietic stem cells. Physicians treat hematological disorders such as anemia, leukemia, and lymphoma through bone marrow transplantation, in which bone marrow is removed from a donor (allogenic transplant) or a patient (autologous transplant) through general surgery, frozen and stored, and then transfused into the patient at a later date. Once transfused into the patient, the bone marrow cells gravitate to the bone marrow and engraft, eventually producing new blood cells either to increase the number of such cells in the anemic patient or to reconstitute the bone marrow destroyed as a result of chemotherapy or radiation therapy. Such methods suffer from the fact that autologous transplants often reintroduced diseased cells into the patient and allogenic transplants caused graft versus host disease.

To avoid such problems, methods have been developed to identify closely-matched donors (with respect to blood and HLA markers) and to isolate the self-renewing hematopoietic stem cells, which generate all red blood cells, neutrophils, T and B lymphocytes, platelets, and macrophages as well as other differentiating cells, from other, diseased, cells. A variety of such separation methods

been developed (see, e.g., PCT patent publication 94/16715), especially for CD34+ cells, a small subset of which are believed to be the ultimate progenitor cell of the hematopoietic system. Due to the problems associated with bone marrow removal surgery and to the advances made in stem cell isolation, however, clinicians developed methods for isolating stem cells from peripheral blood, expanding those cells in culture, and then reintroducing the expanded cells into the patient. During culture of the stem cells, growth factors, such as stem cell factor, G-CSF, erythropoietin, thrombopoietin, IL-3, IL-6, IL-11, and IL-12, are often added to stimulate cell division and differentiation (see, e.g., PCT patent publication No. 94/18991 and EPO patent publication No. 611,263).

Such processes and therapies have been used not only for bone marrow transplantation but also for peripheral blood cell transplantation, replenishment of T cells in AIDS patients, replenishment of lymphocytes in Down's syndrome patients, replenishment of mature immune cells in aplastic anemia patients, and gene therapy. Gene therapy techniques in this context involve the introduction of a genetic construct into explanted cells, expansion of the clones that have incorporated the genetic construct, and then reintroduction of the altered cells into the host (see PCT patent publication Nos. 94/12629, 94/12633, 94/12649, and 94/12650). As methods for isolating cells and differentiating cells in culture continue to improve, the number of disease conditions that can be treated using such methods will increase. For instance, while bone marrow stem cells can be used to regenerate the various differentiated cells of the blood, embryonic and fetal stem cells can be used to regenerate the various differentiated cells of the entire body, offering many diverse applications of cell therapy.

The expansion of cells in culture involves many population doublings, however, so the cells obtained by cell therapy methods after expansion are much closer to senescence than the cells used to begin the process. By practicing the present invention in conjunction with the expansion step, one can prevent the cells from a net loss of

telomeric DNA by lengthening telomeres during the expansion process, while the cells retain the primitive stem cell phenotype and express telomerase activity. The resulting cells are not any closer to senescence than the cells used to begin the process; in fact, the present method offers the possibility that the resulting cells may even have a greater replicative capacity than the cells used to begin the process.

For example, the method of the invention can be applied to bone marrow stem cells, some of which express at least a low level of telomerase activity but have finite replicative capacity. Telomerase activity can be detected in cell-free extracts of dividing, cultured hematopoietic stem and early progenitor cells but not other more differentiated dividing cells. As noted above, numerous kinds of ex vivo cell therapies using bone marrow stem cells are currently practiced or under development, and the present method can readily be practiced in conjunction with such methods. In cases where an individual is going to lose a large population of stem and hematopoietic cells, e.g., chemotherapy or radiation therapy, or is suffering from a disease linked to an immune system undergoing replicative senescence, e.g. normal aging, or cases where the immune system has been severely and chronically stressed, e.g., HIV infection, one can isolate bone marrow stem cells, culture or cultivate those cells in the presence of factors that increase telomere length to slow or reverse replicative senescence (as well as other factors, such as those noted above, if desired), and reintroduce the cells to strengthen or even to reconstitute the immune system.

In the case of AIDS, discussed above, the disease creates a condition in which CD4⁺ cells are limiting, and death is caused by the early senescence of CD4⁺ cells and the resulting unchecked infections by pathogens that follow. It is important to note that such cells age, because the replicative rate of the CD4⁺ cells is increased as a result of HIV infection such that telomere attrition is caused at a greater rate than normal for such cells. The present invention provides therapeutic agents that can be used to treat such diseases and also provides diagnostic procedures by which immune

senescence can be detected and appropriate therapeutic protocols followed.

5 The present invention provides a therapeutic protocol in which CD4⁺ and/or CD8⁺ cells, or a population of cells containing or enriched for CD4⁺ and/or CD8⁺ cells, can be removed from the individual, preferably at an early stage when AIDS is first detected, stored in a bank, and then reintroduced after treatment with a telomere-lengthening reagent of the invention, into the individual at a later stage, 10 i.e., when that individual no longer has a protective amount of CD4⁺ and/or CD8⁺ cells available. Thus, an individual's life can be extended by a protocol involving continued administration of that individual's limiting cells at appropriate time points. These appropriate points can be determined by following 15 CD4⁺ and/or CD8⁺ cell senescence, or by determining the length of telomeres within such CD4⁺ and/or CD8⁺ cells (as an indication of when those cells will become senescent). In the case of AIDS, currently practiced methods include those in which T cells, such as killer T cells, are isolated from a patient and expanded in culture (sometimes after gene therapy to 20 introduce universal receptors for HIV) prior to reintroduction into the patient. The present invention can be used to lengthen the telomeres of such cells either directly or by treatment of a progenitor cells that is then cultured in vitro under differentiating conditions or allowed to differentiate in vivo. 25

Those of skill in the art will recognize that pluripotent hematopoietic stem cells differentiate when grown in culture and that various stimulatory agents, such as cytokines and 30 antigens, can be used to influence the differentiation process. The present method can be practiced in conjunction with such methods to prepare high proliferative capacity cells of a particular type. In similar fashion, embryonic and fetal stem cells are totipotent or pluripotent and can be used to generate 35 cells of any particular type (note that fibroblasts, liver cells, and keratinocytes can all be converted into muscle cells if transfected with vectors that express muscle differentiation factors such as myogenin and myoD). In addition, many late

fetal tissues, such as muscle, lung, kidney, and adrenals, express telomerase activity and so are amenable to the telomere-lengthening methods of the invention. Consequently, the use of totipotent stem cells or telomerase-positive fetal cells in the present method offers significant benefits to the treatment of disease, particularly diseases associated with aging.

Thus, the present invention can be applied to generate cells with high proliferative capacity of a variety of tissue types, including: (a) cells of the central nervous system, including astrocytes, endothelial cells, and fibroblasts, which are involved in such age-related diseases as Alzheimer's disease, Parkinson's disease, Huntington's disease, and stroke; (b) cells of the integument, including fibroblasts, sebaceous gland cells, melanocytes, keratinocytes, Langerhan's cells, and hair follicle cells, which are involved in age-related diseases of the integument, such as dermal atrophy, elastolysis and skin wrinkling, sebaceous gland hyperplasia, senile lentigo, graying of hair and hair loss, chronic skin ulcers, and age-related impairment of wound healing; (c) cells of the articular cartilage, such as chondrocytes, the senescence of which leads to the overexpression of the destructive proteins collagenase and stromelysin, which destroy articular cartilage in osteoarthritis, and lacunal, synovial, connective tissue fibroblasts, which are involved in degenerative joint disease; (d) cells of the bone, such as osteoblasts, osteoclast progenitor cells, bone marrow stromal fibroblasts, and osteoprogenitor cells, which are involved in osteoporosis; (e) cells of the immune system such as B and T lymphocytes, monocytes, neutrophils, eosinophils, basophils, NK cells and their respective progenitors, which are involved in age-related immune system impairment; (f) cells of the vascular system, including endothelial cells, smooth muscle cells, and adventitial fibroblasts, which are involved in age-related diseases of the vascular system including atherosclerosis, calcification, thrombosis, and aneurysms; (g) cells of the eye, such as retinal pigmented epithelium, lens epithelial cells, iris muscle cells (myoblasts), and vascular endothelial cells, which

are involved in loss of vision, i.e., age-related macular degeneration; (h) muscle satellite cells involved in muscular dystrophy; and (i) cells of the gut, such as intestinal epithelial cells, which are involved in malabsorption syndromes.

5 The present invention can therefore be applied to generate a variety of different cell types with high proliferative capacity. Regardless of cell type, practice of the method involves treating the cells with an oligonucleotide substrate for telomerase. In general, such substrates can be
10 identified by their ability to serve as primers for telomerase-mediated nucleotide addition to the 3'-end of the primer in an in vitro reaction. Although the oligonucleotide reagents of the invention can be identified by the property of ability to serve as a substrate for telomerase, the method of the invention in
15 no way requires that the oligonucleotides actually be extended by telomerase in the cell for telomeres to be lengthened. The exact mechanism of action in vivo of the oligonucleotides is not known at this time and may involve titration of a repressor of telomerase or some other
20 mechanism. Indeed, in one embodiment of the invention, no oligonucleotide substrate is required, and the method involves cultivating telomerase-positive cells under conditions where cell division is limited or completely inhibited for a period of time, typically several days to weeks and optionally months,
25 sufficient for telomeres to be lengthened.

Preferably, however, the cells are treated in culture with an oligonucleotide substrate for telomerase in amounts sufficient to lengthen the telomeres of the cells. For human telomerase, such substrates include oligonucleotides that
30 comprise a telomeric repeat sequence 5'-TTAGGG-3'. In particular, oligonucleotides suitable for purposes of the present invention include the 9-, 12-, and 15-nucleotide versions of oligonucleotides that comprise the telomeric repeat sequence, i.e., 5'-TTAGGGTTA-3' and 5'-TTAGGGTTAGGG-3'. Other oligonucleotides that serve as
35 telomerase substrates can also be employed, including the oligonucleotide 5'-TCGAGCACAGTT-3', which corresponds to the junction of a healed broken chromosome from a patient

with thalassemia, and 5'-(GXGXGX)₂₋₃', in which X can be independently selected at each position from either T or A, i.e., 5'-GTGAGTGAGTGA-3'. Other oligonucleotide substrates for telomerase that can be used for purposes of the present invention have been described in the art or can readily be identified using assays described in the art; see, e.g., Morin, 3 October 1991, Nature 353:454-456.

To illustrate how such oligonucleotides are used in the method of the present invention, IDH4, a line of T-antigen immortalized human lung fibroblasts, was cultured in the presence of increasing concentrations of an oligonucleotide containing two telomeric repeats, the GTR oligonucleotide described above. IDH4 cells express telomerase activity and have very short telomeres. Figure 1 shows that the telomeres of IDH4 cells elongate by several kbp after 90 days (about 30 population doublings) of treatment in culture with the GTR oligonucleotide. This elongating effect can be seen both by the increasing size of the DNA resistant to digestion by restriction endonucleases (the "TRF" or "Terminal Restriction Fragment") and by the increasing signal following hybridization to oligonucleotides containing telomeric repeats. The effect is specific to the GTR oligonucleotide, because telomere elongation was not observed following treatment with either an oligonucleotide containing the complementary sequence (5'-CCCTAACCCTAA-3'), called the "CTR" or "C-rich telomeric repeat" oligonucleotide, or a comparably sized oligonucleotide with a randomly selected sequence. The effect may be partially reversible, because IDH4 cells with elongated telomeres exhibited somewhat shorter telomeres when cultured in the absence of oligonucleotides (Figure 1, lane 8, off 90 days). Figure 2 shows that a progressive (up to 30 μ M) dose response is observed with increasing concentrations of the GTR oligonucleotide. Figure 3 shows that the effect of the GTR oligonucleotide is not restricted to a few immortal cell lines, because all of the cell lines tested responded to GTR oligonucleotide treatment by elongating their telomeres. SW620 human colon adenocarcinoma cells, which grow much faster than IDH4 cells, rapidly and progressively elongated

5 their telomeres in the presence of 30 μ M GTR oligonucleotide. The elongation was a continuous process that increased with increasing time of treatment. In addition to SW620 cells (lane 1, control, and lane 2, GTR-treated cells), CAK1 renal carcinoma cells increase the length of their telomeres approximately the same amount as the SW620 GTR-treated cells (Figure 3, lane 3, control, and lane 4, GTR-treated cells). In addition, DU145 prostatic adenocarcinoma cells and COLO205 colon carcinoma cells, which have slightly longer telomeres in control (untreated) cells, also elongate their telomeres in the presence of GTR oligonucleotide (Figure 3, lanes 4-8).

10 While these results were obtained with immortal cells lines to illustrate the invention, the telomere-lengthening effect of the oligonucleotides can be observed in normal cells that express telomerase activity. Methods for measuring telomerase activity have been described in the art (see, e.g., PCT patent publication No. 93/23572), and an especially preferred method is described in copending U.S. patent application Serial No. 08/315,214, filed 28 Sep. 1994 (Harley et al.).

20 Other methods that can be useful in conjunction with the present invention include methods to measure telomere length, which may aid the practitioner in determining whether treatment according to the present method is needed for a particular cell type and whether such treatment has been effective. Procedures for measuring telomere length are known in the art. Typically, restriction endonuclease digestion is used (with enzymes which have a four-base recognition sequence, like HinfI, and do not cleave telomeric DNA) to cleave the genomic DNA; the cleaved fragments are separated according to molecular weight by agarose gel electrophoresis; and the gel is then dried and probed with a radioactively labelled probe, i.e., 5'-(TTAGGG)₃-3', complementary to telomeric DNA sequences.

35 As the results discussed above show, the oligonucleotides of the invention can be transferred into the cytoplasm of a cell spontaneously, i.e., without specific modification. One can

also employ a variety of means to facilitate the uptake of an oligonucleotide by a cell. For instance, one can use liposomes that fuse with the cellular membrane or one can employ ligands that bind to surface membrane protein receptors of the cell resulting in endocytosis of the ligand together with any material bound to the ligand. Alternatively, the cells may be permeabilized to enhance transport of the oligonucleotides into the cell without injuring the cells. One can also use a DNA binding protein, e.g., HBGF-1, which is known to transport an oligonucleotide into a cell.

These delivery modes are useful both for cells in tissue culture (ex vivo) and in vivo. As already indicated, by using liposomes, particularly where the liposome surface carries ligands specific for target cells or are targetted to a specific organ, one can provide for the introduction of an oligonucleotide into the target cells in vivo. For instance, utilizing lipocortin's affinity for phosphatidyl serine, which is released from injured vascular endothelial cells, one can target an oligonucleotide to such a site. Alternatively, catheters, syringes, depots, or the like can be used to provide high localized concentrations of an oligonucleotide in vivo.

In related aspects, the invention features pharmaceutical compositions that include therapeutically effective amounts of the agents described above in pharmaceutically acceptable buffers. These pharmaceutical compositions can include one or more of these agents and be co-administered with other drugs. Compositions and products according to the invention may conveniently be provided in the form of solutions suitable for parenteral, nasal, or oral administration. In many cases, it will be convenient to provide an agent in a single solution for administration.

Amphoteric agents may be utilized as free bases, as acid addition salts, or as metal salts. The salts must, of course, be pharmaceutically acceptable, and these will include metal salts, particularly alkali and alkaline earth metal salts, e.g., potassium or sodium salts. A wide variety of pharmaceutically acceptable acid addition salts are available. These include those prepared from both organic and inorganic

acids, preferably mineral acids. Typical acids that may be mentioned by way of example include citric, succinic, lactic, hydrochloric, and hydrobromic acids. Such products are readily prepared by procedures well known to those skilled in the art.

The agents of the invention can be provided as parenteral compositions for injection or infusion. The agents can, for example, be suspended in an inert oil, suitably a vegetable oil such as sesame, peanut, or olive oil. Alternatively, the agents can be suspended in an aqueous isotonic buffer solution at a pH of about 5.6 to 7.4. Useful buffers include sodium citrate-citric acid and sodium phosphate-phosphoric acid. The desired isotonicity can be achieved using sodium chloride or other pharmaceutically acceptable agents, such as dextrose, boric acid, sodium tartrate, propylene glycol, or other inorganic or organic solutes. Sodium chloride is preferred for buffers containing sodium ions.

If desired, solutions of the above compositions can be thickened with a thickening agent such as methyl cellulose. The solutions can be prepared in emulsified form, either water in oil or oil in water. Any of a wide variety of pharmaceutically acceptable emulsifying agents can be employed, including, for example, acacia powder or an alkali polyether alcohol sulfate or sulfonate.

The therapeutically useful compositions of the invention are prepared by mixing the ingredients following generally accepted procedures. For example, the selected components can be simply mixed in a blender or other standard device to produce a concentrated mixture that can then be adjusted to the final concentration and viscosity by the addition of water or thickening agent and possibly a buffer to control pH or an additional solute to control tonicity.

For use by the physician, the compositions can be provided in dosage unit form containing an amount of agent that will be effective in one or multiple doses to perform the desired function. As will be recognized by those in the field, an effective amount of therapeutic agent will vary with many

factors, including the age and weight of the patient, the patient's physical condition, and other factors.

5 Selected agents, e.g., oligonucleotides and pharmaceutical compositions of the same, can be administered prophylactically or to patients suffering from disease, e.g., by exogenous delivery of the agent to an infected tissue by means of an appropriate delivery vehicle, e.g., a liposome, a controlled release vehicle, by use of iontophoresis, electroporation, or ion-paired molecules, or covalently
10 attached adducts, and other pharmacologically approved methods of delivery. Routes of administration include intramuscular, aerosol, oral (tablet or pill form), topical, systemic, ocular, intraperitoneal, and/or intrathecal. Gene therapy administered using recombinant expression vectors
15 encoding a therapeutic oligonucleotide of the invention can also be used to deliver an oligonucleotide to a cell.

The specific delivery route of any selected agent will depend on the use of the agent. Generally, a specific delivery program for each agent will focus on naked agent uptake with
20 regard to intracellular localization, followed by demonstration of efficacy. Alternatively, delivery to the same cells in an organ or tissue of an animal can be accomplished. Uptake studies can include uptake assays to evaluate cellular oligonucleotide uptake, regardless of the delivery vehicle or
25 strategy. Such assays will also determine the intracellular localization of the agent following uptake, ultimately establishing the requirements for maintenance of steady-state concentrations within the cellular compartment containing the target sequence (nucleus and/or cytoplasm). Efficacy and
30 cytotoxicity can also be tested. Toxicity studies will address not only cell viability but also cell function.

Some methods of delivery suitable for oligonucleotides include: encapsulation in liposomes; transduction by retroviral vectors; and conjugation with agents that direct
35 delivery to certain sites within the body. At least three types of delivery strategies are useful in the present invention, including: agent modification, particle carrier delivery vehicles, and retroviral expression vectors.

Unmodified agents may be taken up by cells slowly and inefficiently. To enhance cellular uptake, an agent can be modified to reduce charge but maintain specific functional groups to improve diffusion across the cell membrane.

5 Modification of agents to reduce charge is one approach to enhance the cellular uptake of these larger molecules. The structural requirements necessary to maintain agent activity are well understood by those in the art. These requirements are taken into consideration when designing modifications to
10 enhance cellular delivery. The modifications are also designed to reduce susceptibility to enzymatic degradation. Both of these characteristics should greatly improve the efficacy of the agent. Chemical modifications of the phosphate backbone of oligonucleotides will reduce the negative charge,
15 allowing free diffusion of the oligonucleotide across the membrane.

 In the body, maintenance of an external concentration is necessary to drive the diffusion of the modified oligonucleotides into the cells of the tissue. Administration
20 routes that allow the diseased tissue to be exposed to a transient high concentration of the oligonucleotide, which is slowly dissipated by systemic adsorption, can be used. Intravenous administration with a drug carrier designed to increase the circulation half-life of the oligonucleotides can
25 also be used. The size and composition of the drug carrier restricts rapid clearance from the blood stream. The carrier, made to accumulate at the site of infection, can protect the oligonucleotides from degradative processes.

 Drug delivery vehicles are effective for both systemic
30 and topical administration. The vehicles can be designed to serve as a slow release reservoir or to deliver their contents directly to the target cell. An advantage of using direct delivery drug vehicles is that multiple molecules are delivered per uptake. Such vehicles have been shown to
35 increase the circulation half-life of drugs that would otherwise be rapidly cleared from the blood stream. Some examples of such specialized drug delivery vehicles that fall into this category are liposomes, hydrogels, cyclodextrins,

biodegradable nanocapsules, and bioadhesive microspheres. For instance, nanoparticles and hydrogels have been used as delivery vehicles for chemotherapeutic agents and protein-based pharmaceuticals.

5 Liposomes increase extracellular stability, increase uptake efficiency and improve biological activity. Liposomes are hollow spherical vesicles composed of lipids arranged in a similar fashion as those lipids that make up the cell membrane. Liposomes have an internal aqueous space for
10 entrapping water soluble compounds and range in size from 0.05 μm to several microns in diameter. Several studies have shown that liposomes can deliver agents to cells in biologically active form. For example, a liposome delivery vehicle originally designed as a research tool, lipofectin, has been
15 shown to deliver intact mRNA molecules to cells yielding production of the corresponding protein. Liposomes offer several advantages: liposomes are non-toxic and biodegradable in composition; liposomes display long circulation half-lives; and recognition molecules can be readily
20 attached to liposome surfaces for targetting to tissues. Finally, cost effective manufacture of liposome-based pharmaceuticals, either in a liquid suspension or lyophilized product, has demonstrated the viability of this technology as an acceptable drug delivery system.

25 Topical administration of agents is advantageous, because topical administration allows localized concentration at the site of administration with minimal systemic adsorption. This simplifies the delivery strategy of the agent to the disease site and reduces the extent of toxicological
30 characterization. Furthermore, the amount of material to be applied is far less than that required for other administration routes. Effective delivery requires the agent to diffuse into the target cells. Chemical modification of the agent to neutralize negative or positive charges can in some instances
35 enhance penetration. The modified agent can also be co-formulated with permeability enhancers, such as azone or oleic acid, in a liposome. The liposomes can either represent a slow release presentation vehicle in which the modified agent

and permeability enhancer transfer from the liposome into the targeted cell, or the liposome phospholipids can participate directly with the modified agent and permeability enhancer in facilitating cellular delivery. In some cases, both the agent and permeability enhancer can be formulated into a suppository formulation for slow release.

Agents can also be systemically administered. Systemic absorption refers to the accumulation of drugs in the blood stream followed by distribution throughout the entire body. Administration routes which lead to systemic absorption include: intravenous, subcutaneous, intraperitoneal, intranasal, intrathecal, and ophthalmic. Each of these administration routes exposes the agent to an accessible diseased or other tissue. Subcutaneous administration drains into a localized lymph node and proceeds through the lymphatic network into the circulation. The rate of entry into the circulation has been shown to be a function of molecular weight and size. The use of a liposome or other drug carrier localizes the agent at the lymph node. The agent can be modified to diffuse into the cell, or the liposome can directly participate in the delivery of either the unmodified or modified agent to the cell.

Thus, delivery methods include liposomes, hydrogels, controlled-release polymers, microinjection, electroporation (for ex vivo treatments), and other pharmaceutically applicable methods. The dosage will depend upon the disease indication and the route of administration but can be between 10-2000 mg/kg of body weight/day. The duration of treatment will extend through the course of the disease symptoms, usually at least 14-16 days and in some indications continuously. Multiple daily doses are anticipated for topical applications, ocular applications, and vaginal applications. The number of doses will depend upon delivery vehicle and efficacy data from clinical trials.

Establishment of therapeutic levels of agent within the target cell is dependent upon the rate of uptake and degradation. Decreasing the degree of degradation will prolong the intracellular half-life of the agent. Thus,

23

chemically modified agents, e.g., oligonucleotides with modification of the phosphate backbone, or capping of the 5' and 3' ends of the oligonucleotides with nucleotide analogues, can be used but may require different dosaging.

5 While the invention is described in a manner that focuses on therapeutic methods for humans and human cells, the method may be applied to cells of any source, particularly cell of animals such as primates and domestic animals such as equine, bovine, avian, ovine, porcine, feline, and canine
10 animals. Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims. The following examples are offered by way of illustration and not by way of limitation.

15 Examples

The following examples illustrate specific aspects of the invention. The examples are not limiting but rather are intended to provide an indication of specific methodology useful in practice of the invention as well as demonstrating
20 the utility of the invention to increase telomere length and slow the onset of cellular senescence. Such correlation indicates to those in the art the breadth of the invention beyond these examples.

25 Example 1

Increasing Telomere Length with GTR Oligonucleotide

An immortalized human fibroblast cell line, IDH4, which has very short TRFs, was incubated with GTR oligonucleotide. Referring to Figs. 1 - 3, cells were incubated in regular culture
30 medium containing serum in the presence of 10 μ M, 30 μ M, and 100 μ M GTR oligonucleotide. The cells were fed fresh phosphodiester GTR oligonucleotide thrice weekly and subcultured when confluent for a total of 90 days. The cells were still growing in GTR oligonucleotide after 90 days at all
35 concentrations used. When TRF analysis was performed after 90 days, the IDH4 cells were shown have increased TRF length in a dose dependent manner with 30 μ M and 100 μ M having approximately the same effect. The control and 30 μ M GTR

oligonucleotide-treated cells were passaged without oligonucleotide addition for an additional 90 days (approximately 35-40 PDL). As is illustrated in Fig. 1, the TRFs slowly shorten.

5 Figure 1 shows that the telomeres of IDH4 cells elongate by several kbp after 90 days (about 30 population doublings) of treatment in culture with the GTR oligonucleotide. This elongating effect can be seen both by the increasing size of the DNA resistant to digestion by restriction endonucleases and by the increasing signal following hybridization to
10 oligonucleotides containing telomeric repeats. The effect is specific to the GTR oligonucleotide, because telomere elongation was not observed following treatment with either an oligonucleotide containing the complementary sequence, or
15 a comparably sized oligonucleotide with a randomly selected sequence. The effect may be partially reversible, because IDH4 cells with elongated telomeres exhibited somewhat shorter telomeres when cultured in the absence of oligonucleotides (Figure 1, lane 8, off 90 days).

20 Figure 2 shows that a progressive (up to 30 μ M) dose response is observed with increasing concentrations of the GTR oligonucleotide. Figure 3 shows that the effect of the GTP oligonucleotide is not restricted to a few immortal cell line because all of the cell lines tested responded to G'
25 oligonucleotide treatment by elongating their telomere. SW620 human colon adenocarcinoma cells, which grow faster than IDH4 cells rapidly and progressively elongate their telomeres in the presence of 30 μ M GTR oligonucleotide. The elongation was a continuous process that increased with increasing time of treatment. In addition to SW620
30 cells (Figure 3, lane 1, control, and lane 2, GTR-treated cells), Caco-2 colon carcinoma cells increase the length of their telomeres approximately the same amount as the SW620 cells (Figure 3, lane 3, control, and lane 4, GTP-treated cells).
35 In addition, DU145 prostatic adenocarcinoma cells and COLO205 colon carcinoma cells, which have telomeres in control (untreated) cells, also

telomeres in the presence of GTR oligonucleotide (Figure 3, lanes 4-8).

Example 2

Increasing Lifespan of Cells

The ability to manipulate telomere length makes possible the demonstration that telomere length determines the lifespan of cells. Cell hybrids were constructed between normal diploid BJ foreskin fibroblasts and either (1) immortal IDH4 cells; or (2) immortal IDH4 cells with telomeres elongated by the method of the present invention. Young BJ cells with approximately 60 doublings left in their normal *in vitro* lifespan were used to ensure that the normal telomeres would not be limiting for the hybrid lifespan.

Fifty clones from each combination were picked and followed to determine their proliferative capacity. The loss of a few normal chromosomes from these hybrids can potentially result in the generation of immortal segregants. To reduce the possibility that a rare segregant would overgrow the population and mask the cessation of growth by the bulk of the cells, the hybrid clones were maintained at low population sizes by serial clonal subcultivation. At each passage, the dishes were scanned, and an average sized clone was picked and replated. Cell counts of parallel dishes of the population of hybrids plated at clonal density indicated that each passage corresponded to approximately 13-14 cell divisions.

More than 95% of the hybrid clones between the immortal IDH4 cells without elongated telomeres and normal BJ fibroblasts ceased proliferating before the third serial subcloning, indicating that such hybrids were only able to undergo approximately 26-39 doublings. In contrast, most of the hybrid clones between the IDH4 cells with elongated telomeres and BJ fibroblasts were able to be maintained for two or more additional passages. The proliferative capacity of the hybrids was thus extended at least an additional 26 doublings by using IDH4 cells whose telomeres were approximately 2 kb longer than the parental IDH4 cells.

To ensure that the above results were not biased by the choice of which subclone was selected for subcultivation, the comparison of the lifespan of the two hybrids was repeated using a protocol designed to determine the lifespan of populations of clones. Hybrids were plated at clonal density in duplicate. After clones had formed, one dish was fixed and counted to determine the number of clones per dish, and one dish was trypsinized and counted to determine the number of cells per dish and then replated to form the next generation of subclones. This permitted the number of cells per clone and thus the average number of cell divisions per passage to be determined. Because dishes containing 100-300 clones were used, the results represent the average of a large number of clones. An additional set of hybrids, comparing parental SW620 cells and SW620 that had been treated with (TTAGG)₂ to elongate its telomeres, was also constructed to verify that the extension of hybrid lifespan was not specific to IDH4 cells. The results showed that the proliferative lifespans of the hybrid cells with elongated telomeres was increased relative to the hybrids with shorter telomeres.

Methods similar to those described in this example can be used to create human hybridoma cell lines with increased proliferative capacity. In this aspect of the invention, an immortal human cell, such as a myeloma cell, is treated according to the method of the present invention to increase telomere length and then fused to a human antibody-producing cell. The resulting cell hybrids have increased proliferative capacity as compared to human hybridoma cells made by methods currently available in the art.

As shown by the above results, by modulating telomere length, one may increase the proliferative capacity of cells and therefore treat disease and disease conditions associated with cellular senescence. All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference. Although the foregoing invention has been described in some detail by way of illustration and

5

example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

Claims

1. A method for increasing the proliferative capacity of normal cells that express telomerase activity, which method comprises culturing or cultivating said cells in the presence of a therapeutically effective amount of an oligonucleotide substrate for telomerase under conditions such that said oligonucleotide substrate enters said cells and acts to lengthen telomeric DNA of said cells.
2. The method of Claim 1, wherein said cells are cultured or cultivated in vivo.
3. The method of Claim 1, wherein said cells are cultured or cultivated ex vivo.
4. The method of Claim 3, wherein said method further comprises delivering cells with lengthened telomeres to a host organism.
5. The method of Claim 1, wherein said cells are embryonic stem cells.
6. The method of Claim 1, wherein said cells are fetal stem cells.
7. The method of Claim 1, wherein said cells are hematopoietic stem cells.
8. The method of Claim 1, wherein said cells are immune system cells.
9. The method of Claim 7, wherein said cells are CD4⁺ cells.
10. The method of Claim 8, wherein said cells are immune system cells of an individual with impaired immune function.

11. The method of Claim 9, wherein said cells are CD4⁺ cells of an HIV-infected individual.

5 12. The method of Claim 1, wherein said oligonucleotide substrate for telomerase is selected from the group of oligonucleotides consisting of oligonucleotides that
10 comprise a telomeric repeat sequence 5'-TTAGGG-3', 5'-TTAGGGTTA-3', 5'-TTAGGGTTAGGG-3', 5'-TCGAGCACAGTT-3',
and 5'-(GXGXGX)₂-3', in which X can be independently selected at each position from either T or A.

15 13. A pharmaceutical composition comprising a pharmaceutically acceptable buffer and a therapeutically effective amount of an agent active to increase length of
telomeres in a cell.

20 14. The pharmaceutical composition of Claim 13, which composition comprises an oligonucleotide substrate for telomerase is selected from the group of oligonucleotides
consisting of oligonucleotides that comprise a telomeric repeat sequence 5'-TTAGGG-3', 5'-TTAGGGTTA-3', 5'-
25 TTAGGGTTAGGG-3', 5'-TCGAGCACAGTT-3', and 5'-(GXGXGX)₂-3', in which X can be independently selected at each position
from either T or A.

15. A method of making a human hybridoma cell with increased proliferative capacity, said method comprising:

- 30 (a) culturing an immortal cell line under conditions such that telomeres in cells of said cell line lengthen; and
 (b) fusing said cells cultured in step (a) with human antibody-producing cells, and said antibody producing cells are human B-lymphocytes.




FIG. 1.




FIG. 2.




FIG. 3.

SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

Internati. Application No
PCT/US 94/13130

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/54 C12N15/10 C12N9/12 C12Q1/48 A61K31/70

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C12Q A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	GENOME, vol. 31, 1989 pages 553-560, BLACKBURN, E. ET AL. 'Recognition and elongation of telomeres by telomerase' cited in the application	1-14
A	see the whole document ---	15
Y	PROC NATL ACAD SCI U S A 89 (21). 1992. 10114-10118, ALLSOPP, R. ET AL. 'TELOMERE LENGTH PREDICTS REPLICATIVE CAPACITY OF HUMAN FIBROBLASTS.' cited in the application see the whole document ---	1-14
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

20 March 1995

Date of mailing of the international search report

04.04.95

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Andres, S

INTERNATIONAL SEARCH REPORT

Internati Application No
PCT/US 94/13130

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MUTAT RES 256 (2-6). 1991. 271-282, HARLEY, C. B. 'TELOMERE LOSS MITOTIC CLOCK OR GENETIC TIME BOMB?.' see the whole document ---	1-15
P,X	WO,A,93 23572 (GERON CORP ;UNIV CALIFORNIA SAN FRANCISCO (US)) 25 November 1993 cited in the application see the whole document ---	1-15
P,X	MOLECULAR AND CELLULAR BIOLOGY 14 (2). 961-969, February 1994 KLINGELHUTZ, A. ET AL. 'Restoration of telomeres in human papillomavirus- immortalized human anogenital epithelial cells.' see the whole document ---	1
T	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA 91 (21). 1994. 9857-9860. ISSN: 0027-8424, VAZIRI H ET AL 'Evidence for a mitotic clock in human hematopoietic stem cells: Loss of telomeric DNA with age.' see the whole document -----	1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 94/ 13130

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 1-12 (as far as in vivo methods are concerned) are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/
composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

Information on patent family members

Internati.

Application No

PCT/US 94/13130

Form PCT/ISA/210 (patent family annex) (July 1992)